# Trends in ADHD Diagnosis

And resultant problems

Bob Pihl

McGill

# Perspective




# A Very Brief History


- 1902 Still- 43 children, symptoms of ADHD, before age 8, 3 to 1 male to female, problems in biological relatives, "cortical disconnection syndrome."
- 1923 Ebaugh et al.- 1917-1918 Encephalitis
   Epidemic , hyperactive, impulsive, attentional-memory deficits. Brain Injury.
- 1930s & 1940s brain injury particular to frontal lobes = ADHD symptoms.

- 1950s- Concept of Minimal Brain Dysfunction.
   Circular but idea brain damage/dysfunction on a continuum. Over stimulated.
- 1960s- Hyperactivity central ( Chess ) DSM II " Hyperkinetic Reaction of Childhood.
- 1970s- Douglas, problem sustaining attention and impulse control. Task performance varies with reinforcement schedule. Attentional problems can persist into later life. DSM III 1980 Attention Deficit Disorder. With and without hyperactivity. Studies suggest under arousal.
- DSM IIIR 1987, Attention Deficit hyperactivity Disorder.

- 1990s to present DSM IV 1994, 3 types, emphasis on genetic, biochemical, brain research.
- Historical Consistencies
- 1. Symptoms fairly consistent- focus on relative importance varied.
  - 2. Gender differences genetic aspects
    - 3. Presumption of brain dysfunction.




Time sitting on a chair plotted against sessions for Cases 1 and 2. After designation A, the patients earned points for remaining seated.




#### Prevalence


- 12-month Prevalence: 4.1% of U.S. adult population<sup>1</sup>
- Severe: 41.3% of these cases (e.g., 1.7% of U.S. adult population) are classified as "severe"<sup>2</sup>




#### **Current ADHD Market Growth is Primarily Driven by Adults**



Source: Age Data - Verispan's Vector One National; Annual Growth - IMS NPA





### For ADHD diagnosis

Combined Type: at least 6 inattention symptoms

Inattentive Type: at least 6 hyperactivity symptoms

Hyperactiveimpulsive Type: at least 6 of each

## ATTENTION DEFICIT HYPERACTIVITY DISORDER <a href="Inattention">Inattention</a>

Fails to give attention to details or makes kareless mistakes

Difficulty sustaining attention in tasks or fun activities

Does not listen when directly spoken to

Does not follow instructions and/or fails to finish work

Has difficulty organizing tasks or activities

Avoids or dislikes tasks requiring a sustained mental effort.

Looses things necessary for tasks or activities

Easily distracted

Forgetful in daily activities

#### **Hyperactivity-Impulsivity**

Fidgets with hands or feet or squirms in seat.

Difficulty remaining seated for prolonged periods

Restless

Difficulty engaging in leisure activities quietly


Always 'on the go' as if 'driven by a motor'


Talks excessively


Blurts out answers before questions have been completed


Difficulty waiting turn

Interrupts or intrudes on others.









#### **Bureaucratic Induced Disorders**

900,000 ADHD's misdiagnosed and treated in the US. (Elder, 2010, Journal of Health Economics)



# Frequency of ADHD Comorbidities

•	Oppositional defiant disorder	40-60%	0/0
•	Conduct disorder	25-35	%
•	Major depressive disorder	30 %	, )
•	Bipolar disorder	10 %	<b>0</b>
•	Anxiety disorder	25 %	, 0
•	Learning disabilities	20 %	<b>/</b> 0

# 33 year follow-up of ADHD's diagnosed at age 8

Significantly worse

Educational, Occupational, economic, and social outcomes.

Ongoing ADHD

Lifetime ASPD & SUD's

More Psychiatric hospitalizations

More Incarcerations, More Divorce

## Evidence of ADHD's relation to DA

- DA medication relatively effective
  - About 80% "successfully "treated
- Animal models of ADHD
  - Frontal lobe lesions of DA projections
    - Producing hyperactivity symptoms
    - Responds to stimulants
- Deficits in DA areas
  - Brain imaging studies: reduced volume in DA rich areas (striatum, frontal lobes)

## Evidence of ADHD's relation to DA

- DA transporter abnormalities
  - Increased DA transporter density in adults with ADHD
  - Results in DA not staying long in synaptic cleft
  - May be related to genetic marker 480 DT allele
    - 3-5 times more likely to have ADHD
- Genetic differences D4 & D2 receptor genes
  - DRD4 gene associated with ADHD
    - 7 repeat allele for this gene increases risk

#### Instruments Used to Assess Externalizing Behavior (EB) and IQ

Table 1. Instruments Used to Assess Externalizing Behavior (EB) and IQ


Sample	Instrument			
Sample 1	EB: Parent and teacher ratings of aggressive and delinquent behaviors on the Child Behavior Checklist and Teacher Report Form. 41 Parent ratings of child and adolescent indicators of psychopathy. 42			
	IQ: WISC-III or WPPSI-R <sup>43,44</sup> ; information and similarities subtests (24 participants); vocabulary and picture completion subtests (16); vocabulary (4); full-scale IQ from prior clinical testing (6).			
Sample 2	EB: Spouse or mother ratings on 3 items: anger control, speech control, and impulsivity level ( $\alpha = .79$ ).  IQ: Full-scale WAIS-III. <sup>45</sup>			
Sample 3	EB: Teacher ratings on aggression, opposition, and hyperactivity scales at age 6 y and yearly from age 10 to 14 y. <sup>1,7,39,40</sup> IQ: WISC-R Vocabulary and Block Design subtests <sup>46</sup> yearly from age 9 to 12 y and at age 15 y; mean number of IQ assessments, 2.47 (SD = 1.67). <sup>40,47</sup>			

Abbreviations: WAIS-III, Wechsler Adult Intelligence Scale, Third Edition; WISC-III, Wechsler Intelligence Scale for Children, Third Edition; WISC-R, Wechsler Intelligence Scale for Children—Revised; WPPSI-R, Wechsler Preschool and Primary Scale of Intelligence—Revised.

DeYoung, C. G. et al. Arch Gen Psychiatry 2006;63:1410-1416.



## Association between externalizing behavior (standardized within each sample) and IQ in 3 male samples, as a function of dopamine D4 receptor 7-repeat allele (DRD4-7) status



DeYoung, C. G. et al. Arch Gen Psychiatry 2006;63:1410-1416.

	DRAFT R	ESEARCH	DOMAIN	CRITERIA	MATRIX	
			UNITS OF ANALYSIS			
DOMAINS/CONSTRUCTS	Genes	Molecules	Cells	Circuits	Behavior	Self-Reports
Negative Affect						
Fear/extinction						
Stress/Distress						
Aggression						
Positive Affect						
Reward Seeking						
Reward/Habit Learning						
Cognition						
Attention						
Perception						
Working Memory						
Declarative memory						
Language behavior						
Cognitive (effortful) control						
Social Processes						
Imitation, theory of mind						
Social dominance						
Facial expression identification						
Attachment/separation fear						
Self-representation areas						
Arousal/Regulatory Processes						
Arousal & regulation (multiple)						
Resting state activity						

## Measuring Impulsivity

Sharma, Markon & Clark (2014)

15 Behavioral Tasks, 4 factors

Inattention, Inhibition, Decision Making, &

Shifting.

14 Self- Report Measures, 3 Factors

Neuroticism, Disinhibition, Extraversion

Low r's between types of measures

Moderate r's to life problems

## Relevant Quotes

" Names are bestowed not found "

"Nosology necessarily precedes etiology"

 "Before I build a wall, I would want to know what I was walling in and walling out"

